



HYPERTHYROIDISM

What is hyperthyroidism?

Hyperthyroidism develops when the body is exposed to excessive amounts of thyroid hormone. This disorder occurs in almost 1 percent of all Americans and affects women five to 10 times more often than men. In its mildest form, hyperthyroidism may not cause recognizable symptoms. However, more often, the symptoms are discomfiting, disabling or even life-threatening.

Features of hyperthyroidism

When hyperthyroidism develops, a goiter (an enlarged thyroid) is usually present and may be associated with some or many of the following features:

- Fast heart rate, often more than 100 beats per minute
- Becoming anxious, irritable, argumentative
- Trembling hands
- Weight loss, despite eating the same amount or even more than usual
- Intolerance of warm temperatures and increased likelihood to perspire
- Loss of scalp hair
- Tendency of fingernails to separate from the nail bed
- Muscle weakness, especially of the upper arms and thighs
- Loose and frequent bowel movements
- Smooth skin
- Change in menstrual pattern
- Increased likelihood for miscarriage
- Prominent “stare” of the eyes
- Protrusion of the eyes, with or without double vision (in patients with Graves’ disease)
- Irregular heart rhythm, especially in patients older than 60 years of age
- Accelerated loss of calcium from bones, which increases the risk of osteoporosis and fractures

Causes of hyperthyroidism

- **Graves’ disease** (named after Irish physician Robert Graves) is an autoimmune disorder that frequently results in thyroid enlargement and hyperthyroidism. In some patients, swelling of the muscles and other tissues around the eyes may develop, causing eye prominence, discomfort or double vision. Like other autoimmune diseases, this condition tends to affect multiple family members. It is much more common in women than in men and tends to occur in younger patients.
- **Toxic multinodular goiter**
Multiple nodules in the thyroid can produce excessive thyroid hormone, causing hyperthyroidism. Typically diagnosed in patients over the age of 50, this disorder is more likely to affect heart rhythm. In many cases, the person has had the goiter for many years before it becomes overactive.
- **Toxic nodule**
A single nodule or lump in the thyroid can produce more thyroid hormone than the body requires and lead to hyperthyroidism. This disorder is not familial.

• Subacute thyroiditis

This condition may follow a viral infection and is characterized by painful enlargement of the thyroid gland and inflammation, which results in the release of large amounts of thyroid hormones into the blood.

Fortunately, this condition usually resolves spontaneously. The thyroid usually heals itself over several months, but often not before a temporary period of low thyroid hormone production (hypothyroidism) occurs.

• Postpartum thyroiditis

Five to 10 percent of women develop mild to moderate hyperthyroidism within several months of giving birth, which typically lasts for one to two months. It is often followed by several months of hypothyroidism, but most women will eventually recover normal thyroid function. In some cases, however, the thyroid gland does not heal and hypothyroidism becomes permanent, requiring lifelong thyroid hormone replacement. Typically, it is not accompanied by a painful thyroid gland. This condition may occur again with subsequent pregnancies.

• Silent thyroiditis

Transient (temporary) hyperthyroidism can be caused by silent thyroiditis, a condition which appears to be the same as postpartum thyroiditis but not related to pregnancy and is not accompanied by a painful thyroid gland.

• Excessive iodine ingestion

Various sources of high iodine concentrations, such as kelp tablets, some expectorants, amiodarone (Cordarone, Pacerone – a medication used to treat heart rhythms) and X-ray dyes, may occasionally cause hyperthyroidism in some patients.

• Overmedication with thyroid hormone

Patients who receive excessive thyroxine replacement treatment can develop hyperthyroidism. They should have their thyroid hormone dosage evaluated by a physician at least once a year and should NEVER take “extra” doses.

Diagnosing hyperthyroidism

Characteristic symptoms and physical signs of hyperthyroidism can be detected by a physician. In addition, tests can be used to confirm the diagnosis and to determine the cause.

• TSH (Thyroid Stimulating Hormone or Thyrotropin) test

A low TSH level in the blood is the most accurate indicator of hyperthyroidism. The body shuts off production of this pituitary hormone when the thyroid gland overproduces thyroid hormone even slightly.

If the TSH level is low, it is important to check thyroid hormone levels as well to confirm the diagnosis of hyperthyroidism.

Continued on back



Other tests

- **Estimates of free thyroxine and free triiodothyronine:**
The active thyroid hormones in the blood. When hyperthyroidism develops, free thyroxine and free triiodothyronine levels rise above previous values – although they may still fall within the normal range for the general population.
- **TSI (Thyroid Stimulating Immunoglobulin):**
A substance often found in the blood when Graves' disease is the cause of hyperthyroidism. This test is ordered if it affects treatment decisions or to help in the diagnosis.
- **Radioactive Iodine Uptake (RAIU):**
A measurement of how much iodine the thyroid gland can collect.
- **Thyroid Scan:**
A thyroid scan shows how the iodine is distributed throughout the thyroid gland. This information can be useful in determining the cause of hyperthyroidism and ultimately its treatment.

Sometimes a general physician can diagnose and treat the cause of hyperthyroidism, but assistance is often needed from an endocrinologist – a physician who specializes in managing thyroid disease.

Treatments

Before the development of current treatment options, the death rate from severe hyperthyroidism was as high as 50 percent. Today, there are several effective treatments and, with proper management, death from hyperthyroidism is rare. Treatment is typically based on the cause of the hyperthyroidism, its severity, and other conditions present. An endocrinologist is a physician experienced in the management of thyroid diseases and who can accurately diagnose the cause of hyperthyroidism and prescribe and manage the best treatment program for each patient.

- **Antithyroid Drugs**
There are two drugs in the U.S. available for the treatment of hyperthyroidism: propylthiouracil (PTU) and methimazole (MMI). In general, the American Association of Clinical Endocrinologists (AACE) and the American Thyroid Association (ATA) recommend prescribing MMI over PTU. However, there are a few situations where PTU should be used over MMI: during the first trimester of pregnancy to avoid an increased risk of a rare birth defect; if the patient is allergic to or intolerant of MMI; or when life-threatening thyrotoxicosis occurs. Some patients with hyperthyroidism caused by Graves' disease experience a spontaneous or natural remission of hyperthyroidism after a 12- to 18-month course of treatment on these drugs and may sometimes avoid permanent underactivity of the thyroid (hypothyroidism), which often occurs as a result of using the other methods of treating hyperthyroidism. Unfortunately, the remission is frequently only temporary, with hyperthyroidism recurring after several months or years off medication and requiring additional treatment. Therefore, relatively few patients are treated solely with antithyroid medication.

Also, antithyroid drugs may cause an allergic reaction in about 5 percent of patients who use them. This usually occurs during the first six weeks of drug treatment. Such a reaction may include rash or hives; but after discontinuing use of the drug, the symptoms resolve within one to two weeks and there is no permanent damage. A more serious effect that occurs in only about 1 in 250-500 patients during the first four to eight weeks of treatment is a rapid decrease of white blood cells in the bloodstream, which may increase susceptibility to serious infection. Symptoms such as a sore throat, infection, or fever should be reported promptly to your physician, and a blood cell count should be done immediately. In most cases, when a person stops using the medication, the white blood cell count returns to normal.

In rare cases, antithyroid drugs may cause severe liver problems, which are detected by blood tests, or joint problems characterized by joint pain and/or swelling. Your physician should be contacted if there is yellowing of the skin ("jaundice"), fever, loss of appetite, or abdominal pain.

Radioactive iodine treatment

Iodine is an essential ingredient in the production of thyroid hormone. Since most overactive thyroid glands are hungry for iodine, it was discovered in the 1940s that the thyroid could be "tricked" into destroying itself by simply feeding it radioactive iodine given by mouth, usually in capsule form which is quickly absorbed from the bowel. It then enters the thyroid cells from the bloodstream and gradually destroys them. Maximal benefit is usually noted within three to six months.

It is not possible to eliminate "just the right amount" of the diseased thyroid gland, since radioiodine eventually damages all thyroid cells. Therefore, most endocrinologists give enough radioactive iodine to make the thyroid underactive (hypothyroidism). Hypothyroidism is easily, predictably and inexpensively corrected by lifelong daily use of oral thyroid hormone replacement therapy.

Although every effort is made to calculate the correct dose of radioiodine for each patient, not every treatment will successfully correct hyperthyroidism, particularly if the goiter is large, and a second dose of radioactive iodine is occasionally needed.

Thousands of patients have received radioiodine treatment, including former U.S. President George H.W. Bush and his wife, Barbara. The treatment appears to be a very safe, simple, and reliably effective one. Because of this, it is considered by most thyroid specialists in the U. S. to be the treatment of choice for hyperthyroidism cases caused by overproduction of thyroid hormone.

Radioactive iodine treatment should never be given to pregnant women or to women who are breastfeeding. Small amounts of radioactive iodine will be excreted in breast milk and could permanently damage the infant's thyroid. Therefore, breastfeeding is not allowed. If radioiodine is inadvertently administered to a woman who is subsequently discovered to be pregnant, the advisability of terminating the pregnancy should be discussed with the patient's OB/GYN and an endocrinologist. Therefore, pregnancy testing is mandatory whenever pregnancy is possible before administering diagnostic or therapeutic radioiodine therapy.

Surgical removal of the thyroid

Though not the preferred treatment for hyperthyroidism, operating to remove most of the thyroid gland may occasionally be recommended in certain situations such as with a pregnant woman with severe uncontrolled disease despite antithyroid drug therapy, since radioiodine would not be safe for the baby. Surgery usually leads to permanent hypothyroidism and lifelong thyroid hormone replacement therapy.

Other treatments

A drug from the class of beta-adrenergic blocking agents (which decrease the effects of excess thyroid hormone) may be used temporarily to control hyperthyroid symptoms until other therapies take effect. In cases where hyperthyroidism is caused by thyroiditis or excessive ingestion of either iodine or thyroid hormone, this may be the only type of treatment required.